


SUSY and the LHC: She was not around the corner

Iacopo Vivarelli

Albert-Ludwigs Universität - Freiburg




Extend the Standard Model?

- The Standard Model is working fine. Why fix it?
 - a) The Higgs sector **suffers from quadratically divergent loop corrections** (high level of fine-tuning)
 - b) **Cosmological data** call for a **dark matter candidate**: no such candidate is present in the SM
 - c) Moreover, **no further unification** between the EW and QCD possible, **no explanation** for EW symmetry breaking, **no easy way** to include gravity, etc.
- Supersymmetry (SUSY) offers an elegant solution for a), which can simultaneously address b) and c)



Supersymmetry (SUSY)




- SUSY is a symmetry that relates bosons and fermions
- In the Minimal Supersymmetric Extension of the Standard Model (MSSM):
 - a new set of fields differing in spin by 1/2 w.r.t. the SM partners (hierarchy problem solved “naturally”)



Mix into neutralinos and charginos

Supersymmetry (SUSY)

- SUSY is a **symmetry that relates bosons and fermions**
- In the Minimal Supersymmetric Extension of the Standard Model (MSSM):
 - a new set of fields differing in spin by 1/2 w.r.t. the SM partners (**hierarchy problem solved “naturally”**)



Mix into
neutralinos and
charginos

$$W \ni \frac{1}{2} \lambda_{ijk} L_i L_j E_k^c + \lambda'_{ijk} L_i Q_j D_k^c + \frac{1}{2} \lambda''_{ijk} U_i^c D_j^c D_k^c + \mu_i L_i H_u$$


- **Lepton and baryon number violation allowed** → proton decay
- If **R-parity conservation assumed**, the **Lightest Supersymmetric Particle (LSP)** is stable: natural Dark Matter candidate

$$R\text{-parity} = (-1)^{3(B-L) + 2s}$$

-1 for sparticles
1 for particles

Why SUSY at the EW scale?



- If SUSY is realised at **some high energy scale** with no measurable effect at the EW scale, **why would we (experimentalist) care?**
- Generic arguments to **expect SUSY to play a role at the EW scale**:
 - **Low fine tuning if $M_{\text{SUSY}} \sim 1 \text{ TeV}$** (therefore within current reach in terms of experiments at colliders)
 - **WIMP miracle of dark matter**:
 - Dark matter relic abundance and interaction cross section related one to the other
 - If cross section EW, then $M_{\text{dark}} \sim 0.1/1 \text{ TeV}$



HEPAP 2006 LHC/ILC Subpanel


SUSY around the corner?

- The Large Hadron Collider has been built to:
 - Fully investigate the mechanism of the EW symmetry breaking**
 - Break the Standard Model:** find SUSY (or any other manifestation of physics beyond the standard model) at the TeV scale
- Early discovery of (EW) R-parity conserving SUSY surely possible by looking at excesses of events with large missing transverse momentum and high jet multiplicity




Prospect for
mSUGRA/CMSSM
discovery (1 fb^{-1})
before switching on
LHC ($\sqrt{s}=10 \text{ TeV}$)

Experimental setup




SUSY and the LHC - 8th January 2013 - LBNL

Experimental setup




Experimental setup




LHC - performance of the machine

- About 22 fb⁻¹ collected at $\sqrt{s} = 8$ TeV and 5 fb⁻¹ at $\sqrt{s} = 7$ TeV (per experiment) in 2011
- Most of which with more than 95% of the detectors operational




Subdetector	Number of Channels	Approximate Operational Fraction
Pixels	80 M	95.0%
SCT Silicon Strips	6.3 M	99.3%
TRT Transition Radiation Tracker	350 k	97.5%
LAr EM Calorimeter	170 k	99.9%
Tile calorimeter	9800	98.3%
Hadronic endcap LAr calorimeter	5600	99.6%
Forward LAr calorimeter	3500	99.8%
LVL1 Calo trigger	7160	100%
LVL1 Muon RPC trigger	370 k	100%
LVL1 Muon TGC trigger	320 k	100%
MDT Muon Drift Tubes	350 k	99.7%
CSC Cathode Strip Chambers	31 k	96.0%
RPC Barrel Muon Chambers	370 k	97.1%
TGC Endcap Muon Chambers	320 k	98.2%

- Most of the analysis so far use 13 fb⁻¹. results being updated for the winter conferences
- **Large luminosity** means **large pileup** (higher than initially planned) . Careful pileup suppression strategies developed.



The detector






- [illegible]

$$M_{eff} = E_T^{miss} + H_T$$

-
- ATLAS Preliminary
- $L^{\text{int}} \sim 35 \text{ pb}^{-1}$
- $H_T^{\mu} [\text{GeV}]$
- Data 2010 ($\sqrt{s} = 7 \text{ TeV}$)
- Standard Model
- MSUGRA $m_0=360 \text{ } m_{1/2}=280$
- Muon Channel
- $E_T^{\text{miss}} > 0.25 m_{\text{eff}}$
- $m_{\text{eff}} > 500 \text{ GeV}$
- $E_T^{\text{miss}} [\text{GeV}]$




Jet measurement

- **Constantly improving** on **jet measurement** and **pileup suppression** techniques
- **Jet energy scale** (isolated jets) known **at the ~1% level**
- thanks to the combination of **several in situ techniques** (gamma/Z+jet balance, di-jet and multi-jet balance, calorimeter/track response comparison)




Missing transverse energy and b-tagging

- Missing E_T (MET) reconstructed from the vectorial sum of **all final state objects**, each with a **dedicated calibration**.
- Large sensitivity of **missing E_T resolution to pileup** addressed by **usage of tracking information in MET calculation**




- **b-tagging:** neural network algorithm combining informations about **secondary vertex displacement** and **impact parameters of jets**
- efficiencies **generally well reproduced by MC**. Systematic uncertainties of the **order of 10-15%**

The Standard Model in one slide





SUSY searches



What processes are we looking for?





What processes are we looking for?



What processes are we looking for?



What processes are we looking for?



- **ATLAS has set up dedicated search strategies for all production mechanism**
- **Only strong production (mainly 3rd generation) covered in this seminar**

Standard Model background estimation


- Doing **SUSY searches** means primarily **understanding the Standard Model background**.
- **Few standard strategies** used all along the ATLAS SUSY analyses (centrally coordinated in the so-called background forum):
 - **Electroweak (and top) production:** rely on fixed order (or NLO) MC generators for predicting shapes of distributions:
 - Normalisation determined in dedicated control regions, shape prediction verified in the control regions and dedicated “validation” regions

Standard Model background estimation

- Doing **SUSY searches** means primarily **understanding the Standard Model background**.
- **Few standard strategies** used all along the ATLAS SUSY analyses (centrally coordinated in the so-called background forum):
 - **Electroweak (and top) production:** rely on fixed order (or NLO) MC generators for predicting shapes of distributions:
 - Normalisation determined in dedicated control regions, shape prediction verified in the control regions and dedicated “validation” regions

• **High cross section processes** can become a relevant **through mechanisms difficult to model with MC:**





- **multi-jet production** is a background for **0-lepton SUSY searches** (through, e.g., fake MET arising from a jet mismeasurement)
- **multi-jet (W+jets)** production is a background for **1-lepton (2-lepton)** analyses if a fake lepton is present in the event
- **Dedicated data driven techniques** used in such cases



Strong SUSY production



Strong production

- Targeting generic strong production of gluinos and squarks.
- The exact decay chain depends on the SUSY mass hierarchy



Strong production

- Targeting generic strong production of gluinos and squarks.
- The exact decay chain depends on the SUSY mass hierarchy



- Two analyses drive the limit with 8 TeV data

	0-lepton (ATLAS-CONF-2012-109)	1-lepton (ATLAS-CONF-2012-104)
leptons	Veto any e or μ above 10 GeV	One isolated e or μ above 25 GeV
jets	2 to 6 jets with $p_T > 60 \text{ GeV}$ (leading jet $p_T > 130 \text{ GeV}$)	4 jets with $p_T > 80 \text{ GeV}$
Other selections	MET > 160 GeV, reject multijet with cuts on MET/ M_{eff} , and angle between jets and MET	MET > 250 GeV, $M_T > 100 \text{ GeV}$, additional selection on MET/ M_{eff}
Final selection	M_{eff}	M_{eff}

Strong production

- **No excess** above SM in any of the signal regions:
 - interpreted **first as a model-independent 95% C.L. limit on σ_{vis}** of BSM processes
 - then as **an exclusion limit in specific SUSY models**

$$\sigma_{\text{vis}} = \sigma \cdot A \cdot \epsilon$$


1-lepton	Signal region	
	Electron	Muon
Observed events	10	4
Fitted background events	9.0 ± 2.8	7.7 ± 3.2
Fitted $t\bar{t}$ events	6.0 ± 2.2	2.6 ± 1.9
Fitted W/Z+jets events	1.5 ± 0.7	4.2 ± 2.3
Fitted other background events	1.0 ± 0.7	0.9 ± 0.3
Fitted multijet events	0.4 ± 0.6	0.0 ± 0.0
MC expected SM events	9.5	11.5
MC expected $t\bar{t}$ events	5.7	4.6
MC expected W/Z+jets events	2.4	6.0
MC expected other background events	1.0	0.8
Data-driven multijet events	0.4	0.0

	$\langle \epsilon \sigma \rangle_{\text{obs}}^{95} [\text{fb}]$	S_{obs}^{95}	S_{exp}^{95}	CL_B
Electron	1.69	9.9	$9.3^{+3.3}_{-2.6}$	0.59
Muon	1.09	6.4	$8.3^{+3.4}_{-2.3}$	0.19

Strong production

- **No excess** above SM in any of the signal regions:
 - interpreted **first as a model-independent 95% C.L. limit on σ_{vis}** of BSM processes
 - then as an **exclusion limit in specific SUSY models**

$$\sigma_{\text{vis}} = \sigma \cdot A \cdot \epsilon$$



1-lepton	Signal region	
	Electron	Muon
Observed events	10	4
Fitted background events	9.0 ± 2.8	7.7 ± 3.2
Fitted $t\bar{t}$ events	6.0 ± 2.2	2.6 ± 1.9
Fitted W/Z+jets events	1.5 ± 0.7	4.2 ± 2.3
Fitted other background events	1.0 ± 0.7	0.9 ± 0.3
Fitted multijet events	0.4 ± 0.6	0.0 ± 0.0
MC expected SM events	9.5	11.5
MC expected $t\bar{t}$ events	5.7	4.6
MC expected W/Z+jets events	2.4	6.0
MC expected other background events	1.0	0.8
Data-driven multijet events	0.4	0.0

	$\langle \epsilon \sigma \rangle_{\text{obs}}^{95} [\text{fb}]$	S_{obs}^{95}	S_{exp}^{95}	CL_B
Electron	1.69	9.9	$9.3^{+3.3}_{-2.6}$	0.59
Muon	1.09	6.4	$8.3^{+3.4}_{-2.3}$	0.19

- Simplified models: **assume degenerate 1st and 2nd generation squarks**. The only possible production processes are $gg, qq, q\bar{q}$
- only possible processes (depending on masses)

$$\tilde{g} \rightarrow qq\tilde{\chi}_1^0, \tilde{g} \rightarrow \tilde{q}\tilde{\chi}_1^0, \tilde{q} \rightarrow q\tilde{\chi}_1^0$$


Squark (gluino) masses below 1.3/1.4 (1.1) TeV excluded for any gluino (squark) mass

Natural SUSY

- **She was not around the corner. Is she then gone?**
 - Or, more specifically: is the argument of SUSY solving the SM fine tuning not an argument anymore?

M.Papucci, J.Ruderman, A. Weiler

- “Naturalness” in SUSY driven by **the μ parameter, the stop mass** (the gluino mass to a lesser extent)
- “Light” stop(s), light sbottom (left), lighter higgsinos, not-so-heavy gluinos
- Almost no constraint on the other sparticles
- The general MSSM **can accommodate** $m_h = 125/126$ GeV and keep low fine tuning (see, for example M. W. Cahill-Rowley et al., arXiv:1206.5800[hep-ph])





Gluino mediated stop production

- Gluinos and stops/sbottoms the only “light” strong interacting particles:
 - **gluino mediated stop/sbottom production**
 - **direct pair production of stops/sbottoms**

Gluino mediated stop production

- Gluinos and stops/sbottoms the only “light” strong interacting particles:

- **gluino mediated stop/sbottom production**
- **direct pair production of stops/sbottoms**



- If gluino pair production dominant (and only stops not too heavy), then the decay is

$$\tilde{g} \rightarrow \tilde{t}t$$

- Final state that contains **MET (LSP)**, up to 4 b-jets, up to 12 jets, up to 4 leptons (possibly same sign)
- Three different analyses target this final state:
 - **3-b jets plus MET** (up to 6 jets) - ATLAS-CONF-2012-145
 - **2 SS leptons + MET + 4 jets** - ATLAS-CONF-2012-105
 - **multijet (up to 9 jets)** - ATLAS-CONF-2012-103
 - **3-leptons + MET** - ATLAS-CONF-2012-151

direct 3rd generation squark production

- The **stops (and the sbottom left)** constrained by naturalness to be **not heavier than ~ 1 TeV**
- Wide, dedicated effort for **direct stop production search** in ATLAS


- The t_1 decay branching ratios depend strongly on:

- Mass hierarchy ($\tilde{t}_1, \tilde{X}_1^0, \tilde{X}_1^\pm$)
- Chirality of the stop and of the $\tilde{X}_1^0, \tilde{X}_1^\pm$


- Several analyses used:

- $\tilde{t}_1 \rightarrow t\tilde{X}_1^0$: 0-lepton, 1-lepton*, 2-leptons
- $\tilde{t}_1 \rightarrow b\tilde{X}_1^\pm$: 1-lepton, 2-leptons*, 0-lepton (2b + MET)*

- * = analyses updated for $\sqrt{s} = 8$ TeV or new




2011 status



2 b-jets + MET


- Generic topology of many interesting physics channels:
 - In particular sbottom and stop pair production



2 b-jets + MET


- Generic topology of many interesting physics channels:
 - In particular sbottom and stop pair production

- If the chargino and neutralino are degenerate in mass, the final states are identical \Rightarrow **target small $\Delta m(\tilde{X}_1^\pm, \tilde{X}_1^0)$**
- Basically two signal regions used



2 b-jets + MET

- Generic topology of many interesting physics channels:
 - In particular sbottom and stop pair production




- If the chargino and neutralino are degenerate in mass, the final states are identical \Rightarrow **target small $\Delta m(\tilde{X}_1^\pm, \tilde{X}_1^0)$**
- Basically two signal regions used

- m_{CT} : boost-corrected contranverse mass
 - $m_{CT}^2(b_1, b_2) = (E_T(b_1) + E_T(b_2))^2 - (\mathbf{p}_T(b_1) - \mathbf{p}_T(b_2))^2$
 - for pair produced particles decaying into visible and invisible particles
 - It has an end-point at $(m_{prod}^2 - m_{inv}^2)/m_{prod}^2$

- **SR1(for large $\Delta m(\tilde{b}/\tilde{t}, \tilde{X}_1^0)$):**
 - Look for 2 b-jets (veto on third jet), large MET
 - Use M_{CT} to suppress top (end-point at 135 GeV)
 - Main background (at high M_{CT}): $Z (\rightarrow \nu\nu) + b$ -jets
 - SR2 is similar

2 b-jets + MET

- Generic topology of many interesting physics channels:
 - In particular sbottom and stop pair production



- If the chargino and neutralino are degenerate in mass, the final states are identical \Rightarrow **target small $\Delta m(\tilde{X}_1^\pm, \tilde{X}_1^0)$**
- Basically two signal regions used

- m_{CT} : boost-corrected contranverse mass
 - $m_{CT}^2(b_1, b_2) = (E_T(b_1) + E_T(b_2))^2 - (\mathbf{p}_T(b_1) - \mathbf{p}_T(b_2))^2$
 - for pair produced particles decaying into visible and invisible particles
 - It has an end-point at $(m_{prod}^2 - m_{inv}^2)/m_{prod}^2$

- **SR1 (for large $\Delta m(\tilde{b}/\tilde{t}, \tilde{X}_1^0)$):**
 - Look for 2 b-jets (veto on third jet), large MET
 - Use M_{CT} to suppress top (end-point at 135 GeV)
 - Main background (at high M_{CT}): $Z (\rightarrow \nu\nu) + b$ -jets
 - SR2 is similar

- **SR3 (for small $\Delta m(\tilde{b}/\tilde{t}, \tilde{X}_1^0)$):**
 - Focus on events with a hard ISR jet produced
 - Hard, non b-tagged leading jet, two additional b-jets
 - Veto on additional hadronic activity
 - Main background: top pair production


Control region definition

- Background estimation

Control region definition

• Background estimation

- top/W control region:
 - 2 b-jets
 - 1-lepton, $40 \text{ GeV} < M_T < 100 \text{ GeV}$, similar selection as SR
 - At high M_{CT} W and top both relevant



Control region definition



• Background estimation

• top/W control region:

- 2 b-jets
- 1-lepton, $40 \text{ GeV} < M_T < 100 \text{ GeV}$, similar selection as SR
- At high M_{CT} W and top both relevant

• Z control region:

- 2 b-jets
- 2-lepton same flavour, select the Z peak
- “Mimic” MET by “neutrinising” the leptons



Control region definition

• Background estimation

• top/W control region:




- 2 b-jets
- 1-lepton, $40 \text{ GeV} < M_T < 100 \text{ GeV}$, similar selection as SR
- At high M_{CT} W and top both relevant

• Z control region:

- 2 b-jets
- 2-lepton same flavour, select the Z peak
- “Mimic” MET by “neutrinoising” the leptons

• top control region:

- 2 b-jets
- 2-lepton different flavour
- Very pure top control region



Control region counts and systematics

ATLAS-CONF-2012-165


Channel	CR1L_SR1	CR2L_SR1	CR2LDF_SR1
Observed events	104	102	51
Fitted bkg events	104 ± 11	102 ± 11	51 ± 7
Top production	70 ± 16	18 ± 4	50 ± 7
Z production	1.5 ± 0.4	82 ± 12	—
W production	25 ± 19	—	—
Others	8 ± 4	2.4 ± 1.3	0.8 ± 0.4

- Systematic uncertainties:

- b-tagging uncertainties (~15%)
- jet energy scale uncertainty (~10%)
- Z production theoretical uncertainties (5%)

• Results **compatible with SM background predictions** in all signal regions

- Normalisation factors for the backgrounds in control regions close to 1 for top, to 1.2 for Z



Channel	SR1, m_{CT} selection				SR2	SR3	
	150 GeV	200 GeV	250 GeV	300 GeV		SR3a	SR3b
Observed	172	66	16	8	104	207	21
SM Total	176 ± 25	71 ± 11	25 ± 4	7.4 ± 1.7	95 ± 11	203 ± 35	27 ± 5
Top production	45 ± 13	17 ± 6	7 ± 3	1.6 ± 0.6	15 ± 4	146 ± 40	15 ± 5
Z production	85 ± 15	36 ± 6	12 ± 2	4.0 ± 0.9	60 ± 9	27 ± 9	7 ± 2
W production	28 ± 23	12 ± 10	4 ± 3	1 ± 1	15 ± 5	22 ± 7	4 ± 1
Others	6 ± 3	4 ± 2	1.4 ± 0.8	0.7 ± 0.4	4 ± 2	4 ± 2	1.5 ± 0.9
Multijet production	12 ± 12	2 ± 2	0.2 ± 0.2	0.01 ± 0.01	0.6 ± 0.6	4 ± 4	—

0-lepton, 2-b jets stop searches

- 95% C.L. model independent upper limits on BSM event yield and σ_{vis}

Signal region	Bkg. estimate	Obs. data	95% CL UL on BSM event yield		95% CL UL on σ_{vis} (fb)	
			expected	observed	expected	observed
SR1 ($m_{\text{CT}} > 150$ GeV)	176 ± 25	172	55	54	4.2	4.1
SR1 ($m_{\text{CT}} > 200$ GeV)	71 ± 11	66	25	22	1.9	1.7
SR1 ($m_{\text{CT}} > 250$ GeV)	25 ± 4	16	12.5	7.9	0.96	0.61
SR1 ($m_{\text{CT}} > 300$ GeV)	7.4 ± 1.7	8	7.5	8.0	0.58	0.62
SR2	95 ± 11	104	32	39	2.5	3.0
SR3a	203 ± 35	207	54	54	4.2	4.2
SR3b	27 ± 5	21	13.1	9.6	1.0	0.74


0-lepton, 2-b jets stop searches

- 95% C.L. model independent upper limits on BSM event yield and σ_{vis}

Signal region	Bkg. estimate	Obs. data	95% CL UL on BSM event yield		95% CL UL on σ_{vis} (fb)	
			expected	observed	expected	observed
SR1 ($m_{\text{CT}} > 150$ GeV)	176 ± 25	172	55	54	4.2	4.1
SR1 ($m_{\text{CT}} > 200$ GeV)	71 ± 11	66	25	22	1.9	1.7
SR1 ($m_{\text{CT}} > 250$ GeV)	25 ± 4	16	12.5	7.9	0.96	0.61
SR1 ($m_{\text{CT}} > 300$ GeV)	7.4 ± 1.7	8	7.5	8.0	0.58	0.62
SR2	95 ± 11	104	32	39	2.5	3.0
SR3a	203 ± 35	207	54	54	4.2	4.2
SR3b	27 ± 5	21	13.1	9.6	1.0	0.74



- 95% C.L. limit **very similar** for the sbottom and the stop case if $\Delta m(\tilde{X}_1^\pm, \tilde{X}_1^0) = 5$ GeV
- A clear **loss of acceptance** (because of lepton and jet veto) if $\Delta m(\tilde{X}_1^\pm, \tilde{X}_1^0) = 20$ GeV

- $\tilde{t} \rightarrow b \tilde{X}_1^\pm$ (BR 100%) excluded up to $m_t \sim 600$ GeV for nearly degenerate chargino and neutralino masses**




1-lepton stop searches (ATLAS-CONF-2012-166)


- Dedicated **1-lepton analysis** dominates the reach for $t\bar{t}$ pair production ($\tilde{t}_1 \rightarrow \tilde{t} X_1^0$)
- Basic strategy: **select top-like events with large M_T and MET.**
- Main background: **$t\bar{t} \rightarrow l\bar{l}bb + \text{MET}$ (dominant at large M_T)**
- Dedicated effort to **improve the second lepton veto** (isolated tracks - tau-like events)



Direct stop search summary





- **Mixed decays** (50% $\tilde{t}_1 \rightarrow t \tilde{X}_1^0$, 50% $\tilde{t}_1 \rightarrow b \tilde{X}_1^\pm$) still not considered (and somewhat favoured, actually)
- Complete the investigation in **the low mass region** (exclude independently of the stop parameters and mass hierarchy).





Prospects for SUSY searches at 13/14 TeV

- LHC is **foreseen to run at 13/14 TeV** after 2015 and **integrate about 300 fb⁻¹**.
- Increased **energy and pileup conditions** (highly dependent on the bunch spacing)
 - Impact mainly on trigger conditions:
 - Short term: improve on trigger strategy - trigger on topologies
 - Long term: dedicated hardware/software upgrades: improved calorimeter readout at early trigger stages, single track trigger, etc.
- Expect to deal with **increased p_T thresholds** at the beginning, especially for leptons (lowest unprescaled lepton trigger of $p_T > 33$ GeV w.r.t. current 25 GeV)



Prospects with 13/14 TeV

- Project the sensitivity of the analyses to **13/14 TeV**
- ...assuming **realistic running conditions** and no improvement on the analyses (!)
- A lot still to be said about EW scale SUSY



Summary

- **SUSY was not around the corner.** Degenerate **1st and 2nd generation squarks (gluinos)** are excluded up to **$m = 1.5 \text{ TeV}$ (1.2 TeV)** for $m_{\text{LSP}} = 0$.
- Searches for direct production **of third generation squarks** are also very well advanced.
 - **Stops decaying (only) into top-LSP** excluded up to **$\sim 560 \text{ GeV}$** for massless LSP. Large fraction of the available parameter space also excluded for $\tilde{t} \rightarrow b \tilde{X}_1^\pm$
 - The 14 TeV data will allow to probe up to $m_{\text{stop}} \sim 1 \text{ TeV}$
- The **startup of the LHC** has been a fruitful and exciting period for SUSY searches
 - Waiting for the 13/14 TeV running


BACKUP

1-lepton stop searches (ATLAS-CONF-2012-166)

- Dedicated **1-lepton analysis** dominates the reach for t_1 pair production ($t_1 \rightarrow tX_1^0$)
- Basic strategy: **select top-like events with large M_T and MET.**
- Main background: **$t\bar{t} \rightarrow l\bar{l}b\bar{b}$ + MET (dominant at large M_T)**
- Dedicated effort to **improve the second lepton veto** (isolated tracks - tau-like events)




Select m_{jjj} compatible with had top decay

Requirement	SRD	SRE	SRtN1	SRtN2	SRtN3	SRbC
$\Delta\phi(j_1, \vec{p}_T^{\text{miss}}) >$	0.8	0.8	0.8	-	0.8	0.8
$\Delta\phi(j_2, \vec{p}_T^{\text{miss}}) >$	0.8	0.8	0.8	0.8	0.8	0.8
$E_T^{\text{miss}} [\text{GeV}] >$	225	275	150	200	225	150
$E_T^{\text{miss}} / \sqrt{H_T} [\text{GeV}^{1/2}] >$	11	11	8	13	11	7
$m_T [\text{GeV}] >$	130	140	140	140	180	120
$m_T [\text{GeV}] <$	-	-	250	-	-	-
$am_{T2} [\text{GeV}] >$	-	-	-	170	200	-
$m_{T2}^{\tau} [\text{GeV}] >$	-	-	-	-	120	-
$N^{\text{iso-trk}} = 0$	-	-	-	-	-	Yes
$A \times \varepsilon$ benchmark point	-	-	0.06%	0.9%	2.8%	0.7%



1-lepton stop searches

- **top pair production background** normalised in control region ($60 \text{ GeV} < M_T < 90 \text{ GeV}$). **Same for W** (anti-b-tag)
- Fake lepton background obtained with Matrix Method
- No excess in any of the signal regions.
- Result interpreted in $t_1 t_1$ production with $t_1 \rightarrow t X_1^0$ (BR 100%) or $t_1 \rightarrow b X_1^\pm$ (BR 100%)



Signal region definition

SR1:

- ▶ 2 b -jets (130, 50) GeV
- ▶ veto 3rd jet (above 50 GeV)
- ▶ $m_{CT} > 150, 200, 250, 300$ GeV

SR2:

- ▶ 2 b -jets (60, 60)
- ▶ veto 3rd jet (above 50 GeV)
- ▶ $E_T^{\text{miss}} > 200$ GeV
- ▶ $m_{CT} > 100$ GeV
- ▶ $HT,2 < 50$ GeV

$$HT,2 = \sum_{\text{jet} > 2}^{\text{jet } N} p_T$$

SR3 (a):



- ▶ ISR jet 130 GeV, anti b -tagged
- ▶ 2 b -jets (30, 30) GeV
- ▶ $\Delta\Phi(E_T^{\text{miss}}, \text{jet1}) > 2.5$
- ▶ $p_T \text{ } b\text{-jet } 1 < 110$ GeV
- ▶ $HT,3 < 50$ GeV

SR3 (b):

- ▶ all SR3(a) cuts
- ▶ $E_T^{\text{miss}} > 250$
- ▶ $p_T \text{ jet1} > 150$ GeV


Missing transverse energy

- Missing E_T (MET) reconstructed from the vectorial sum of **all final state objects**, each **with a dedicated calibration**.
- Large sensitivity of **missing E_T resolution to pileup** addressed by **usage of tracking information in MET calculation**




Fake MET background estimate

- Large MET can be induced by a jet mis-measurement. Relevant for processes with high cross section and no “real” MET (multi-jet, $Z \rightarrow \ell\ell$)





- Derive a “jet response function” from the MC and adapt it to reproduce:

- p_T balance in di-jet events (core of response function)
- response tail in three-jet (Mercedes) events



- Use response function to smear real data events with low MET:
- Obtain events with large “fake” MET
- Validate them in a dedicated control region (MET aligned with one jet)



Fake lepton background estimate


- General approach to **fake lepton background estimation** based on a **loose/tight matrix method**
- Example with 1 lepton (easily extendable to multi-lepton signatures): **multijet events can fake 1-lepton signatures** if:
 - a lepton from a **heavy flavour hadron decay** passes the lepton selection
 - electrons from **photon conversions** pass the lepton selection
- Strategy: **define a “loose”** (pre-selected) **and a “tight”** (signal) lepton selection.
- Then, simply solve the following system of equations

$$\begin{aligned}
 N^{\text{loose}} &= N_{\text{real}}^{\text{loose}} + N_{\text{fake}}^{\text{loose}} \\
 N^{\text{tight}} &= \epsilon_{\text{real}} N_{\text{real}}^{\text{loose}} + \epsilon_{\text{fake}} N_{\text{fake}}^{\text{loose}}
 \end{aligned}$$

Need to be measured independently



Simply count how many of them

$$N_{\text{fake}}^{\text{tight}} = \frac{\epsilon_{\text{fake}}}{\epsilon_{\text{real}} - \epsilon_{\text{fake}}} (N_{\text{real}}^{\text{loose}} \epsilon_{\text{real}} - N^{\text{tight}})$$



gluino mediated stop production

- Gluinos and stops/sbottoms the only “light” strong interacting particles:
 - gluino mediated stop/sbottom production
 - direct pair production of stops/sbottoms





- If gluino pair production dominant (and only stops not too heavy), then the decay is $\tilde{g} \rightarrow \tilde{t}t$
- Final state that contains MET (LSP), up to 4 b-jets, up to 12 jets, up to 4 leptons (possibly same sign)
- Three different analyses target this final state:
 - 3-b jets plus MET (up to 6 jets) - ATLAS-CONF-2012-145
 - 2 SS leptons + MET + 4 jets - ATLAS-CONF-2012-105
 - multijet (up to 9 jets) - ATLAS-CONF-2012-103
 - 3-leptons + MET - ATLAS-CONF-2012-151

Strong production

- No excess above SM in any of the signal regions:
 - interpreted first as a model-independent 95% C.L. limit on the visible cross section of BSM processes
 - then as an exclusion limit in specific SUSY models

1-lepton	Electron	Muon
Observed events	10	4
Fitted background events	9.0 ± 2.8	7.7 ± 3.2
Fitted $t\bar{t}$ events	6.0 ± 2.2	2.6 ± 1.9
Fitted W/Z+jets events	1.5 ± 0.7	4.2 ± 2.3
Fitted other background events	1.0 ± 0.7	0.9 ± 0.3
Fitted multijet events	0.4 ± 0.6	0.0 ± 0.0
MC expected SM events	9.5	11.5
MC expected $t\bar{t}$ events	5.7	4.6
MC expected W/Z+jets events	2.4	6.0
MC expected other background events	1.0	0.8
Data-driven multijet events	0.4	0.0

1-lepton	$\langle \epsilon \sigma \rangle_{\text{obs}}^{95} [\text{fb}]$	S_{obs}^{95}	S_{exp}^{95}	CL_B
Electron	1.69	9.9	$9.3^{+3.3}_{-2.6}$	0.59
Muon	1.09	6.4	$8.3^{+3.4}_{-2.3}$	0.19




Strong production


- Simplified models often used in the limit setting
 - Only a few SUSY particles play a role, all others are set to very high masses
- In this case, assume degenerate 1st and 2nd generation squarks. The only possible production processes are

$$\tilde{g}\tilde{g}, \tilde{g}\tilde{q}, \tilde{q}\tilde{q}$$
- only possible processes (depending on masses)

$$\tilde{g} \rightarrow qq\tilde{\chi}_1^0, \tilde{g} \rightarrow \tilde{q}\tilde{\chi}_1^0, \tilde{q} \rightarrow q\tilde{\chi}_1^0$$
- Squark (gluino) masses below 1.3/1.4 (1.1) TeV excluded for any gluino (squark) mass



Strong production




Main SM background processes: top pair production, W production


- Addressed for both analyses with 1-lepton control regions (by selecting the M_T Jacobian peak)
- W and top regions separated by the use of b-tagging

- $Z \rightarrow \nu\nu$ (for 0-lepton) addressed by using a photon + jet control region
- Multijet background addressed using the jet smearing method (0-lepton) or the matrix method (1-lepton)

$Z \rightarrow \nu\nu$ control region




top control region





2-leptons stop searches

- A dedicated two-lepton analysis addresses best the case $t_1 \rightarrow bX_1^\pm$ (BR 100% and $X_1^\pm \rightarrow W^\pm X_1^0$) if $\Delta m(t_1, X_1^\pm)$ not too small
- Same and different flavour leptons considered
- Main SM background: tt , WZ
- Basic variable used to reject tt , WW : M_{T2}

$$m_{T2}(\mathbf{p}_T^{\ell_1}, \mathbf{p}_T^{\ell_2}, \mathbf{p}_T^{\text{miss}}) = \min_{\mathbf{q}_T + \mathbf{r}_T = \mathbf{p}_T^{\text{miss}}} \left\{ \max[m_T(\mathbf{p}_T^{\ell_1}, \mathbf{q}_T), m_T(\mathbf{p}_T^{\ell_2}, \mathbf{r}_T)] \right\}$$




The exclusion limit concentrated in regions with small $\Delta m(t_1, X_1^\pm)$




Is then SUSY dead?

- **No!** (pretty consistent theoretical view on the subject)
- Gluinos and 1st and 2nd generation squarks can be heavy, provided that the **superpartners of the heavy fermions are relatively light** (still “natural” hierarchy)
- Higgsinos should also be light



Most exciting, alive + natural SUSY spectrum




N. Arkani-Ahmed (CERN Workshop, 31-Oct'11)

... and this is not yet covered by LHC

3rd generation production

- The third generation squarks are special
- **Mixing between L and R component** of squarks proportional to the fermion partner mass → **large mixing for 3rd generation squarks** → generally expected to be **the lightest squarks**
- Previous limits on squark masses do not apply to sbottom and stop
- The stop directly counter-balance quadratically divergent top loop corrections to the Higgs mass



0-lepton results

Signal Region	A-loose	A-medium	B-medium	C-loose	C-medium	E-loose	E-medium
MC expected events							
Diboson	53.1	18.2	11.1	6.2	0.9	0.0	0.0
W+jets	264.1	53.5	51.9	62.9	16.4	2.1	1.9
Z/ γ^* +jets	338.2	74.7	50.4	55.0	16.1	1.0	0.8
$t\bar{t}$ + single top	74.9	8.1	14.2	42.6	5.3	2.1	1.6
Fitted background events							
Diboson	53 \pm 23	18 \pm 9	11 \pm 6	6 \pm 4	0.9 \pm 0.6	–	–
Multi-jets	0.6 \pm 0.6	0.1 \pm 0.1	0.2 \pm 0.2	–	–	–	–
W+jets	180 \pm 140	33 \pm 35	32 \pm 34	40 \pm 40	8 \pm 8	1.2 \pm 1.3	0.9 \pm 1.1
Z/ γ^* +jets	354 \pm 21	81 \pm 8	59 \pm 6	67 \pm 6	18.5 \pm 3.0	2.0 \pm 1.0	0.6 \pm 0.5
$t\bar{t}$ + single top	67 \pm 16	7.6 \pm 3.5	14 \pm 5	39 \pm 7	5.3 \pm 2.0	2.5 \pm 0.9	2.0 \pm 1.4
Total bkg	650 \pm 130	140 \pm 33	115 \pm 30	155 \pm 31	33 \pm 8	5.7 \pm 1.7	3.5 \pm 1.7
Observed	643	111	106	156	31	9	7
p_0	0.498	0.500	0.500	0.486	0.498	0.161	0.108
UL on N_{BSM}	224.8	33.9	43.8	65.7	17.9	10.4	9.9
UL on σ_{BSM} (fb)	38.8	5.84	7.55	11.3	3.09	1.79	1.71

Signal Region	A-tight	B-tight	C-tight	D-tight	E-tight
MC expected events					
Diboson	3.3	0.2	0.0	0.8	2.6
W+jets	6.6	5.6	2.1	3.4	3.3
Z/ γ^* +jets	7.4	4.5	1.9	1.3	1.3
$t\bar{t}$ + single top	1.0	1.1	0.6	1.8	2.7
Fitted background events					
Diboson	3.3 \pm 3.1	0.2 \pm 1.4	–	0.8 \pm 0.4	2.6 \pm 2.0
Multi-jets	–	–	–	0.4 \pm 0.5	0.1 \pm 0.2
W+jets	3 \pm 4	2.7 \pm 3.4	0.3 \pm 0.5	–	0.8 \pm 1.3
Z/ γ^* +jets	6.8 \pm 2.2	5.1 \pm 1.7	2.0 \pm 1.1	2.5 \pm 1.1	1.2 \pm 0.7
$t\bar{t}$ + single top	0.8 \pm 0.8	0.8 \pm 0.9	0.6 \pm 0.5	2.6 \pm 1.6	5.1 \pm 3.3
Total bkg	14 \pm 5	8.7 \pm 3.4	2.8 \pm 1.2	6.3 \pm 2.1	10 \pm 4
Observed	10	7	1	5	9
p_0	0.499	0.500	0.499	0.500	0.499
UL on N_{BSM}	8.9	7.3	3.3	6.0	9.3
UL on σ_{BSM} (fb)	1.53	1.26	0.57	1.03	1.60

CR	SR background	CR process	CR selection
CRY	Z($\rightarrow \nu\nu$)+jets	γ +jets	Isolated photon
CRQ	QCD jets	QCD jets	Reversed $\Delta\phi(\text{jet}, E_T^{\text{miss}})_{\text{min}}$ and $E_T^{\text{miss}}/m_{\text{eff}}(Nj)$ cuts
CRW	W($\rightarrow \ell\nu$)+jets	W($\rightarrow \ell\nu$)+jets	30 GeV < $m_T(\ell, E_T^{\text{miss}})$ < 100 GeV, b -veto
CRT	$t\bar{t}$ and single- t	$t\bar{t} \rightarrow b\bar{b}q\bar{q}'\ell\nu$	30 GeV < $m_T(\ell, E_T^{\text{miss}})$ < 100 GeV, b -tag

Minimal SUSY extension of SM (MSSM)

- If SUSY is unbroken, $M_{\text{particle}} = M_{\text{particle}}$. Since sparticles are not yet observed, **SUSY must be (softly) broken**:

- $\mathcal{L}_{\text{SUSY}} = \mathcal{L}_{\text{SUSY conserving}} + \mathcal{L}_{\text{SUSY soft breaking}}$

2 higgs doublets
needed \rightarrow 5 Higgs bosons

MSSM parameters:

SUSY conserving sector	SUSY breaking sector
3 coupling constants for SU(3) x SU(2)xSU(1)	5 3x3 hermitian mass matrices (one per EW multiplet)
4 Yukawa couplings per generation	3 complex 3x3 matrices (Higgs trilinear couplings to sfermions)
	3 mass terms for the Higgs sector + 2 additional off-diagonal terms
	Higgs VEV expectation angle β

A total of 124 parameters.



But: separate lepton number conservation, FCNC suppression, CP violation leave about 20 independent parameters

Alternatively: Precise assumptions can be made on the way SUSY is broken:

- mSUGRA/CMSSM
- GMSB
- AMSB

(R-parity) conserving SUSY phenomenology



- Missing E_T **is the main signature** (although not unique)
- General R-parity signatures:
 - The LSP (typically $\tilde{\chi}_1^0$ or $\tilde{\nu}$ in mSUGRA) is stable and weakly interacting (dark matter candidate) \rightarrow **large missing transverse momentum**
 - squarks and gluinos produced in pp collision \rightarrow **large particle multiplicity** typically produced in the decay.
- Large jet multiplicities and/or lepton produced in large regions of the parameter space (although not mandatory, e.g. $pp \rightarrow \tilde{q}\tilde{q} \rightarrow qq \tilde{\chi}_1^0 \tilde{\chi}_1^0$)



RPV stop searches

$$W \ni \frac{1}{2} \lambda_{ijk} L_i L_j E_k^c + \lambda'_{ijk} L_i Q_j D_k^c + \frac{1}{2} \lambda''_{ijk} U_i^c D_j^c D_k^c + \mu_i L_i H_u$$

- Depending on which couplings differ from zero, one can get different signatures:
 - Highly ionising tracks (R-hadrons)
 - Two-jet resonances
 - Lot of work still to be done here




Higgs and SUSY

$$X_t = (A_t + \mu \cot \beta) / m_S$$


$$m_h^2 = m_Z^2 \cos^2 \beta + \frac{3y_t^2 m_t^2}{(4\pi)^2} \left[\log \left(\frac{m_S^2}{m_t^2} \right) + X_t^2 \left(1 - \frac{X_t^2}{12} \right) \right]$$

arXiv:1212.6847

- The Higgs mass depend on the average stop mass and X_t
- $m_h=126$ GeV still allows for a light t_1



ATLAS in a nutshell (II)



A dileptonic $t\bar{t}b\bar{b}$ event candidate

An ATLAS event

